

INTEGRALLY CLOSED IDEALS ON LOG TERMINAL SURFACES ARE MULTIPLIER IDEALS

KEVIN TUCKER

ABSTRACT. We show that all integrally closed ideals on log terminal surfaces are multiplier ideals by extending an existing proof for smooth surfaces.

CONTENTS

1. Introduction	1
2. Background	2
2.1. Relative Numerical Decomposition	2
2.2. Antinef Closures and Global Sections	2
2.3. Generic Sequences of Blowups	3
3. Main Theorem	4
3.1. Log Terminal Singularities and Multiplier Ideals	4
3.2. Choosing $\mathfrak{a}$ and λ	4
3.3. Conclusion of Proof	5
References	7

1. INTRODUCTION

Throughout this article, we will consider a scheme $X = \operatorname{Spec} \mathcal{O}_X$, where $\mathcal{O}_X$ is a two-dimensional local normal domain essentially of finite type over $\mathbb{C}$. Our purpose is to address the following question, raised in [LLS08]:

Question. If X has a rational singularity, is every integrally closed ideal a multiplier ideal?

When X is regular, an affirmative answer was given concurrently by [LW03] and [FJ05]. Our main result is to generalize their methods to prove the following:

Theorem 1.1. *Suppose X has log terminal singularities. Then every integrally closed ideal is a multiplier ideal.*

Recall that log terminal singularities are necessarily rational (see Theorem 5.22 in [KM98]). In a sense, this gives a complete answer to the above question: if X has a rational singularity but fails to be log terminal, every multiplier ideal is strictly proper. In particular, $\mathcal{O}_X$ itself is not a multiplier ideal.

The author was partially supported by the NSF under grant DMS-0502170.

There are several difficulties in trying to extend the techniques used in [LW03]. One must show that successful choices can be made in the construction (specifically, the choice of ϵ and N in Lemma 2.2 of [LW03]). Here, it is essential that X has log terminal singularities. Further problems arise from the failure of unique factorization to hold for integrally closed ideals. As X is not necessarily factorial, we may no longer reduce to the finite colength case. In addition, the crucial contradiction argument which concludes the proof in [LW03] does not apply. These nontrivial difficulties are overcome by using a relative numerical decomposition for divisors on a resolution over X .

Our presentation is self-contained and elementary. Section 2 contains background material covering the relative numerical decomposition, antinef closures, and some computations using generic sequences of blowups. Section 3 is dedicated to the constructions and arguments in the proof of Theorem 1.1.

2. BACKGROUND

2.1. Relative Numerical Decomposition. Consider $X = \text{Spec } \mathcal{O}_X$, where $\mathcal{O}_X$ is a two-dimensional local normal domain essentially of finite type over $\mathbb{C}$. Let $x \in X$ be the unique closed point, and suppose $f : Y \rightarrow X$ is a projective birational morphism such that Y is regular and $f^{-1}(x)$ is a simple normal crossing divisor. Let $E_1, \dots, E_u$ be the irreducible components of $f^{-1}(x)$, and $\Lambda = \oplus_i \mathbb{Z}E_i \subset \text{Div}(Y)$ the lattice they generate.

The intersection pairing $\text{Div}(Y) \times \Lambda \rightarrow \mathbb{Z}$ induces a negative definite $\mathbb{Q}$ -bilinear form on $\Lambda_{\mathbb{Q}}$ (see [Art66] for an elementary proof). Consequently, there is a dual basis $\check{E}_1, \dots, \check{E}_u$ for $\Lambda_{\mathbb{Q}}$ defined by the property that

$$\check{E}_i \cdot E_j = -\delta_{ij} = \begin{cases} -1 & i = j \\ 0 & i \neq j \end{cases}.$$

Recall that a divisor $D \in \text{Div}_{\mathbb{Q}}(Y)$ is said to be f -antinef if $D \cdot E_i \leq 0$ for all $i = 1, \dots, u$. In this case, D is effective if and only if f_*D is effective (see Lemma 3.39 in [KM98]). In particular, $\check{E}_1, \dots, \check{E}_u$ are effective.

If $C \in \text{Div}_{\mathbb{Q}}(X)$, we define the numerical pullback of C to be the unique $\mathbb{Q}$ -divisor f^*C on Y such that $f_*f^*C = C$ and $f^*C \cdot E_i = 0$ for all $i = 1, \dots, u$. Note that, when C is Cartier or even $\mathbb{Q}$ -Cartier, this agrees with the standard pullback of C . If $D \in \text{Div}_{\mathbb{Q}}(Y)$, we have

$$(1) \quad D = f^*f_*D + \sum_i (-D \cdot E_i) \check{E}_i.$$

We shall refer to this as a relative numerical decomposition for D . Note that, even when D is integral, both f^*f_*D and $\check{E}_1, \dots, \check{E}_u$ are likely non-integral. The fact that f^*f_*D and $\check{E}_1, \dots, \check{E}_u$ are always integral divisors when X is smooth and D is integral is equivalent to the unique factorization of integrally closed ideals. See [Lip69] for further discussion.

2.2. Antinef Closures and Global Sections. Suppose now that $D' = \sum_E a'_E E$ and $D'' = \sum_E a''_E E$ are f -antinef divisors, where the sums range over the prime divisors E on Y . It is easy

to check that $D' \wedge D'' = \sum_E \min\{a'_E, a''_E\}E$ is also f -antinef. Further, any integral $D \in \text{Div}(Y)$ is dominated by some integral f -antinef divisor (e.g. $(f^{-1})_* f_* D + M(\check{E}_1 + \cdots + \check{E}_u)$ for sufficiently large and divisible M). In particular, there is a unique smallest integral f -antinef divisor $D^\sim$, called the f -antinef closure of D , such that $D^\sim \geq D$. One can verify that $f_* D = f_* D^\sim$, and in addition the following important lemma holds (see Lemma 1.2 of [LW03]). The proof also gives an effective algorithm for computing f -antinef closures.

Lemma 2.1. *For any $D \in \text{Div}(Y)$, we have $f_* \mathcal{O}_Y(-D) = f_* \mathcal{O}_Y(-D^\sim)$.*

Proof. Let $s_D \in \mathbb{N}$ be the sum of the coefficients of $D^\sim - D$ when written in terms of $E_1, \dots, E_n$. If $s_D = 0$, then $D = D^\sim$ is f -antinef and the statement follows trivially. Else, there is an index i such that $D \cdot E_i > 0$. As $E_i \cdot E_j \geq 0$ for $j \neq i$, we must have

$$D \leq D + E_i \leq D^\sim = (D + E_i)^\sim.$$

Thus, $s_{D+E_i} = s_D - 1$ and by induction we may assume

$$f_* \mathcal{O}_Y(-(D + E_i)) = f_* \mathcal{O}_Y(-(D + E_i)^\sim) = f_* \mathcal{O}_Y(-D^\sim)$$

and it is enough to show $f_* \mathcal{O}_Y(-D) = f_* \mathcal{O}_Y(-(D + E_i))$. Consider the exact sequence

$$0 \longrightarrow \mathcal{O}_Y(-(D + E_i)) \longrightarrow \mathcal{O}_Y(-D) \longrightarrow \mathcal{O}_{E_i}(-D) \longrightarrow 0.$$

Since $\deg(\mathcal{O}_{E_i}(-D)) = -D \cdot E_i < 0$, we have $f_* \mathcal{O}_{E_i}(-D) = 0$; applying f_* yields the desired result. $\square$

2.3. Generic Sequences of Blowups. In the proof of Theorem 1.1, we will make use of the following auxiliary construction. Suppose $x^{(i)}$ is a closed point of E_i with $x^{(i)} \notin E_j$ for $j \neq i$. A generic sequence of n -blowups over $x^{(i)}$ is:

$$Y = Y_0 \xleftarrow{\sigma_1} Y_1 \xleftarrow{\sigma_2} \cdots \xleftarrow{\sigma_{n-1}} Y_{n-1} \xleftarrow{\sigma_n} Y_n$$

where $\sigma_1 : Y_1 \rightarrow Y_0$ is the blowup of $Y_0 = Y$ at $x_1 := x^{(i)}$, and $\sigma_k : Y_k \rightarrow Y_{k-1}$ is the blowup of Y_{k-1} at a generic closed point x_k of $(\sigma_{k-1})^{-1}(x_{k-1})$ for $k = 2, \dots, n$. Let $\sigma : Y_n \rightarrow Y$ be the composition $\sigma_n \circ \cdots \circ \sigma_1$. We will denote by $E(1), \dots, E(u)$ the strict transforms of $E_1, \dots, E_u$ on Y_n . Also, let $E(i, x^{(i)}, k)$, $k = 1, \dots, n$, be the strict transforms of the n new σ -exceptional divisors created by the blowups $\sigma_1, \dots, \sigma_n$, respectively.

Lemma 2.2. (a.) *Let $\sigma : Y_n \rightarrow Y$ be a generic sequence of blowups over $x^{(i)} \in E_i$. Then one has*

$$\check{E}(i) \leq \check{E}(i, x^{(i)}, 1) \leq \cdots \leq \check{E}(i, x^{(i)}, n).$$

(b.) *Suppose $D \in \text{Div}(Y)$ is an integral $(f \circ \sigma)$ -antnef divisor such that E_i is the unique component of $\sigma_* D$ containing $x^{(i)}$. If $\text{ord}_{E(i)} D = a_0$ and $\text{ord}_{E(i, x^{(i)}, k)} D = a_k$ for $k = 1, \dots, n$, then*

$$a_0 \leq a_1 \leq \cdots \leq a_n.$$

Further, $a_0 < a_n$ if and only if

$$\left(\sum_{k=1}^n (-D \cdot E(i, x^{(i)}, k)) \check{E}(i, x^{(i)}, k) \right) \geq \check{E}(i).$$

Proof. If $n = 1$, we have

$$\begin{aligned} \check{E}(i, x^{(i)}, 1) &= \left(\sigma^* \check{E}_i + E(i, x^{(i)}, 1) \right) \geq \sigma^* \check{E}_i = \check{E}(i) \\ D &= \sigma^* \sigma_* D + (-D \cdot E(i, x^{(i)}, 1)) \check{E}(i, x^{(i)}, 1). \end{aligned}$$

The general case of both statements follows easily by induction. $\square$

3. MAIN THEOREM

3.1. Log Terminal Singularities and Multiplier Ideals. Once more, suppose $x \in X$ is the unique closed point and $f : Y \rightarrow X$ is a projective birational morphism such that Y is regular and $f^{-1}(x)$ is a simple normal crossing divisor. Let $E_1, \dots, E_u$ be the irreducible components of $f^{-1}(x)$, and let K_Y be a canonical divisor on Y . Then $K_X := f_* K_Y$ is a canonical divisor on X . If we write the relative canonical divisor as

$$K_f := K_Y - f^* K_X = \sum_i b_i E_i$$

then X has log terminal singularities if and only if $b_i > -1$ for all $i = 1, \dots, u$. In this case, X is automatically $\mathbb{Q}$ -factorial (see Proposition 4.11 in [KM98], as well as [dFH08] for recent developments).

If $\mathfrak{a} \subseteq \mathcal{O}$ is an ideal, recall that $f : Y \rightarrow X$ as above is said to be a log resolution of $\mathfrak{a}$ if $\mathfrak{a}\mathcal{O}_Y = \mathcal{O}_Y(-G)$ for an effective divisor G such that $\text{Ex}(f) \cup \text{Supp}(G)$ has simple normal crossings. In this case, we can define the multiplier ideal of $(X, \mathfrak{a})$ with coefficient $\lambda \in \mathbb{Q}_{>0}$ as

$$\mathcal{J}(X, \mathfrak{a}^\lambda) = f_* \mathcal{O}_Y(\lceil K_f - \lambda G \rceil).$$

See [Tuc07] for an introduction in a similar setting, or [Laz04] for a more comprehensive overview. Also recall that $\mathfrak{a}$ is integrally closed if and only if

$$\mathfrak{a} = f_* \mathcal{O}_Y(-G).$$

3.2. Choosing $\mathfrak{a}$ and λ . We now begin the proof of Theorem 1.1. For the remainder, assume X is log terminal, and let $I \subseteq \mathcal{O}_X$ be an integrally closed ideal. In this section, we construct another ideal $\mathfrak{a} \subseteq \mathcal{O}_X$ along with a coefficient $\lambda \in \mathbb{Q}_{>0}$, and in the following section it will be shown that $\mathcal{J}(X, \mathfrak{a}^\lambda) = I$. Let $f : Y \rightarrow X$ a log resolution of I with exceptional divisors $E_1, \dots, E_u$. Suppose $I\mathcal{O}_Y = \mathcal{O}_Y(-F^0)$, and write

$$K_f = \sum_{i=1}^u b_i E_i$$

$$F^0 = (f^{-1})_* f_* (F^0) + \sum_{i=1}^u a_i E_i.$$

Choose $0 < \epsilon < 1/2$ such that $\lfloor \epsilon(f^{-1})_* f_* (F^0) \rfloor = 0$ and

$$\epsilon(a_i + 1) < 1 + b_i$$

for $i = 1, \dots, u$. Note that, since X is log terminal, $1 + b_i > 0$ and any sufficiently small $\epsilon > 0$ will do. Let $n_i := \lfloor \frac{1+b_i}{\epsilon} - (a_i + 1) \rfloor \geq 0$, and $e_i := (-F^0 \cdot E_i)$. Choose e_i closed points $x_1^{(i)}, \dots, x_{e_i}^{(i)}$ on E_i such that $x_j^{(i)} \notin \text{Supp}((f^{-1})_* f_* (F^0))$ and $x_j^{(i)} \notin E_l$ for $l \neq i$. Denote by $g : Z \rightarrow Y$ the composition of n_i generic blowups at each of the points $x_j^{(i)}$ for $j = 1, \dots, e_i$ and $i = 1, \dots, u$. As in Section 2.3, denote by $E(1), \dots, E(u)$ the strict transforms of $E_1, \dots, E_u$, and $E(i, x_j^{(i)}, 1), \dots, E(i, x_j^{(i)}, n_i)$ the strict transforms of the n_i exceptional divisors over $x_j^{(i)}$.

Let $h := f \circ g$, $F = g^*(F^0)$, and choose an effective h -exceptional integral divisor A on Z such that $-A$ is h -ample. It is easy to see that

$$K_g = \sum_{i=1}^u \sum_{j=1}^{e_i} \sum_{k=1}^{n_i} k E(i, x_j^{(i)}, k)$$

and one checks

$$K_g \cdot E(i) = e_i \quad K_g \cdot E(i, x_j^{(i)}, k) = \begin{cases} 0 & k \neq n_i \\ -1 & k = n_i \end{cases}.$$

It follows immediately that $F + K_g$ is h -antinef. Choose $\mu > 0$ sufficiently small that

$$(2) \quad \lfloor (1 + \epsilon)(F + K_g + \mu A) - K_h \rfloor = \lfloor (1 + \epsilon)(F + K_g) - K_h \rfloor.$$

As $-(F + K_g + \mu A)$ is h -ample, there exists $N \gg 0$ such that $G := N(F + K_g + \mu A)$ is integral and $-G$ is relatively globally generated.¹ In other words, $\mathfrak{a} := h_* \mathcal{O}_Z(-G)$ is an integrally closed ideal such that $\mathfrak{a} \mathcal{O}_Z = \mathcal{O}_Z(-G)$. Set $\lambda = \frac{1+\epsilon}{N}$.

3.3. Conclusion of Proof. Here, we will show $\mathcal{J}(X, \mathfrak{a}^\lambda) = I = h_* \mathcal{O}_Z(-F)$. Since

$$\mathcal{J}(X, \mathfrak{a}^\lambda) = h_* \mathcal{O}_Z(\lceil K_h - \lambda G \rceil) = h_* \mathcal{O}_Z(-\lfloor \lambda G - K_h \rfloor),$$

by Lemma 2.1, it suffices to show $F' := \lfloor \lambda G - K_h \rfloor^\sim = F$. In particular, we have reduced to showing a purely numerical statement.

Lemma 3.1. *We have $F' \leq F$ and $h_* F' = h_* F$. In addition, for $i = 1, \dots, u$ and $j = 1, \dots, e_i$,*

$$\text{ord}_{E(i, x_j^{(i)}, n_i)}(F') = \text{ord}_{E(i, x_j^{(i)}, n_i)}(F) = \text{ord}_{E(i)}(F).$$

¹As X is log terminal, it also has rational singularities, and by Theorem 12.1 of [Lip69] it follows that $-(F + K_g)$ is already globally generated without the addition of $-A$. However, the above approach seems more elementary, and avoids unnecessary reference to these nontrivial results.

Proof. Since $F' = [\lambda G - K_h]^\sim$ and F is h -antinef ($-F$ is relatively globally generated), it suffices to show these statements with $[\lambda G - K_h]$ in place of F' . By (2), we have

$$\begin{aligned} [\lambda G - K_h] &= \lfloor (1 + \epsilon)(F + K_g) - K_h \rfloor \\ &= F + \lfloor \epsilon(F + K_g) - g^*K_f \rfloor. \end{aligned}$$

Since $\lfloor \epsilon(f^{-1})_* f_* F^0 \rfloor = 0$, it follows immediately that $h_*[\lambda G - K_h] = h_*F$. For the remaining two statements, consider the coefficients of $\epsilon(F + K_g) - g^*K_f$. Along $E(i)$, we have $\epsilon a_i - b_i$, which is less than one by choice of ϵ . Along $E(i, x_j^{(i)}, k)$, we have $\epsilon(a_i + k) - b_i$. This expression is greatest when $k = n_i$, where our choice of n_i guarantees

$$\begin{aligned} \frac{b_i}{\epsilon} - a_i &\leq n_i < \frac{b_i + 1}{\epsilon} - a_i \\ 0 &\leq \epsilon(a_i + n_i) - b_i < 1. \end{aligned}$$

It follows that $[\lambda G - K_h] \leq F$, with equality along $E(i, x_j^{(i)}, n_i)$. $\square$

Lemma 3.2. *For each $i = 1, \dots, u$,*

$$(-F' \cdot E(i))\check{E}(i) + \sum_{j=1}^{e_i} \sum_{k=1}^{n_i} (-F' \cdot E(i, x_j^{(i)}, k))\check{E}(i, x_j^{(i)}, k) \geq (-F \cdot E(i))\check{E}(i).$$

Proof. If $\text{ord}_{E(i)} F' = \text{ord}_{E(i)} F$, as $F' \leq F$ we have $F' \cdot E(i) \leq F \cdot E(i)$ and the conclusion follows as $\check{E}(i)$ and $\check{E}(i, x_j^{(i)}, k)$ are effective and F' is h -antinef. Otherwise, if $\text{ord}_{E(i)} F' < \text{ord}_{E(i)} F = \text{ord}_{E(i, x_j^{(i)}, n_i)} F'$, then for each $j = 1, \dots, e_i$ we saw in Lemma 2.2(b) that

$$\sum_{k=1}^{n_i} (-F' \cdot E(i, x_j^{(i)}, k))\check{E}(i, x_j^{(i)}, k) \geq \check{E}(i).$$

Summing over all j gives the desired conclusion. $\square$

We now finish the proof by showing that $F' \geq F$. Using the relative numerical decomposition (1) and the previous two Lemmas, we compute

$$\begin{aligned} F' &= h^* h_* F' + \sum_{i=1}^u (-F' \cdot E(i))\check{E}(i) + \sum_{i=1}^u \sum_{j=1}^{e_i} \sum_{k=1}^{n_i} (-F' \cdot E(i, x_j^{(i)}, k))\check{E}(i, x_j^{(i)}, k) \\ &= h^*(h_* F) + \sum_{i=1}^u \left((-F' \cdot E(i))\check{E}(i) + \sum_{j=1}^{e_i} \sum_{k=1}^{n_i} (-F' \cdot E(i, x_j^{(i)}, k))\check{E}(i, x_j^{(i)}, k) \right) \\ &\geq h^* h_* F + \sum_{i=1}^u (-F \cdot E(i))\check{E}(i) = F. \end{aligned}$$

This concludes the proof of Theorem 1.1.

REFERENCES

- [Art66] Michael Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136. MR MR0199191 (33 #7340)
- [dFH08] Tommaso de Fernex and Christopher D. Hacon, *Singularities on normal varieties*, arXiv:0805.1767, 2008.
- [FJ05] Charles Favre and Mattias Jonsson, *Valuations and multiplier ideals*, J. Amer. Math. Soc. **18** (2005), no. 3, 655–684 (electronic). MR MR2138140 (2007b:14004)
- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. MR MR1658959 (2000b:14018)
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. II*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 49, Springer-Verlag, Berlin, 2004, Positivity for vector bundles, and multiplier ideals. MR MR2095472 (2005k:14001b)
- [Lip69] Joseph Lipman, *Rational singularities, with applications to algebraic surfaces and unique factorization*, Inst. Hautes Études Sci. Publ. Math. (1969), no. 36, 195–279. MR MR0276239 (43 #1986)
- [LLS08] Robert Lazarsfeld, Kyungyong Lee, and Karen E. Smith, *Syzygies of multiplier ideals on singular varieties*, Michigan Mathematics Journal (2008).
- [LW03] Joseph Lipman and Kei-ichi Watanabe, *Integrally closed ideals in two-dimensional regular local rings are multiplier ideals*, Math. Res. Lett. **10** (2003), no. 4, 423–434. MR MR1995782 (2004m:13059)
- [Tuc07] Kevin Tucker, *Jumping numbers on algebraic surfaces with rational singularities*, To appear. arXiv:0801.0734v2, 2007.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109

E-mail address: kevtuck@umich.edu